

A GIAMBELLI FORMULA FOR ISOTROPIC PARTIAL FLAG VARIETIES

HARRY TAMVAKIS

ABSTRACT. Let G be a classical complex Lie group, P any parabolic subgroup of G , and G/P the corresponding partial flag variety. We prove an explicit nonrecursive Giambelli formula which expresses an arbitrary Schubert class in $H^*(G/P)$ as a polynomial in certain special Schubert class generators; in the even orthogonal case this depends on the main result of [BKT4]. Our formula extends to one that applies to the torus-equivariant cohomology ring of G/P and to the setting of symplectic and orthogonal degeneracy loci.

0. INTRODUCTION

The classical Giambelli formula [G] is one of the fundamental results concerning Schubert calculus in the cohomology ring of the usual Grassmannian $\mathfrak{X}$. The cohomology of $\mathfrak{X}$ has an additive basis of Schubert classes, and is generated as a ring by certain *special* Schubert classes, which are the Chern classes of the universal quotient bundle over $\mathfrak{X}$. The formula of Giambelli writes a general Schubert class as a determinant of a Jacobi-Trudi matrix with entries given by special classes.

The Giambelli problem can be formulated for any homogeneous space G/P , where G is a classical complex Lie group and P a parabolic subgroup of G . One can choose as Schubert class generators for the cohomology ring $H^*(G/P)$ the pullbacks of the special Schubert classes on Grassmannians; typically these will be the Chern classes of the tautological quotient bundles over G/P . The question then is to find an explicit combinatorial formula which expresses a general Schubert class in $H^*(G/P)$ as a polynomial in the special classes. We proceed to give a brief survey of the work to date on the Giambelli problem for such homogeneous spaces.

The case of the full flag variety G/B , where the cohomology is generated by Schubert divisors, is more amenable to study. Note that in general, the Giambelli polynomials are only defined up to the ideal of relations generated by the invariant polynomials under the action of the Weyl group of G . Bernstein-Gelfand-Gelfand [BGG] and Demazure [D1, D2] used divided difference operators to construct an algorithm that produces polynomials which represent the Schubert classes. For the general linear group, Lascoux and Schützenberger [LS1] applied this method to define *Schubert polynomials*, a particularly nice choice of representatives with rich combinatorial properties. An explicit combinatorial formula for the coefficients of Schubert polynomials was given by Billey, Jockusch, and Stanley [BJS] – thus resolving the Giambelli question in the case of the complete flag variety in type A.

Date: August 25, 2009.

2000 *Mathematics Subject Classification.* Primary 14M15; Secondary 05E15, 14M17, 14N15, 05E05.

The author was supported in part by NSF Grant DMS-0901341.

This was generalized to arbitrary GL_n partial flag varieties by Buch, Kresch, Yong, and the author [BKTY, §5].

In the other classical Lie types, one considers the varieties which parametrize flags of subspaces of a vector space which are isotropic with respect to an orthogonal or symplectic form. Pragacz [P] found an explicit Giambelli formula for the maximal isotropic Grassmannians, in the form of a Schur Pfaffian [Sch]. At the other extreme, for the full flag varieties G/B in types B, C, and D, the analogous theory of Schubert polynomials is not uniquely determined (see [BH, Fu2, FK2, LP1, LP2, T1, T3] for examples) and their combinatorics is more challenging.

Giambelli formulas for non-maximal isotropic Grassmannians were discovered only recently [BKT2, BKT4]. Unlike most previously known examples, the nature of the formulas in loc. cit. is not determinantal – instead, they are expressed using Young’s raising operators [Y]. In the present work we will generalize these results to settle the Giambelli problem for any classical G/P space (when G is an even orthogonal group, the problem is reduced to the main result of [BKT4]). One of our motivations for this is the fact that all the known Giambelli formulas expressed in terms of special Schubert classes have straightforward – often identical – extensions to the small quantum cohomology ring of G/P ; see [Be, CF, FGP, KT2, KT3, BKT3]. We expect that our new Giambelli formulas will similarly prove to be the right ones to look at in the quantum world.

The modern formulation of the Giambelli problem is in the setting of an algebraic family of varying partial flag varieties, with applications to *degeneracy loci* of vector bundles. This story also has a long history, from the work of Thom-Porteous and Kempf-Laksov [KL] to the generalizations by Fulton, Pragacz, and others [Fu1, Fu2, Fu3, Fu4, PR, LP1, KT1, IMN]; see [FP, Ma, Tu] for expositions. It has been known at least since the paper of Graham [Gra] that the degeneracy locus problem for the classical groups from [Fu2, Fu3] is equivalent to the Giambelli problem for the Schubert classes in the T -equivariant cohomology of G/P , where T denotes the maximal torus in G . In type A, the project culminated with the combinatorial understanding of the polynomials representing quiver loci [BF, BKTY, KMS].

Fulton asked in [Fu3] for degeneracy locus formulas which have a similar shape for all the classical groups; moreover, that the formulas should be determinantal whenever possible. We give a complete answer to this question here, except that in addition to determinants we also require the raising operator Giambelli polynomials from [BKT2, BKT4]. Furthermore, the results provide a combinatorial link between the general G/P Giambelli problem and the quiver formulas of [BKTY].

We now state one of our main theorems, referring to §1, §3, and §5 for the precise definitions. Equip the vector space $E = \mathbb{C}^{2n}$ with a nondegenerate skew-symmetric bilinear form. Fix a sequence $\mathfrak{d} : d_1 < \cdots < d_p$ of positive integers with $d_p \leq n$. Let $\mathfrak{X}(\mathfrak{d})$ be the variety parametrizing partial flags of subspaces

$$(1) \quad E_{\bullet} : 0 \subset E_1 \subset \cdots \subset E_p \subset E$$

with $\dim E_i = d_i$ for each i and E_p isotropic. The Schubert varieties $\mathfrak{X}_w$ in $\mathfrak{X}(\mathfrak{d})$ and their cohomology classes $[\mathfrak{X}_w]$ are indexed by signed permutations w in the Weyl group of Sp_{2n} whose descent positions are included among the d_i . Let $E'_j = E/E_{p+1-j}$ for $1 \leq j \leq p$; the Chern classes of the E'_j are the special Schubert

classes in $H^*(\mathfrak{X}(\mathfrak{d}), \mathbb{Z})$. We then have

$$(2) \quad [\mathfrak{X}_w] = \sum_{\underline{\lambda}} e_{\underline{\lambda}}^w \Theta_{\lambda^1}(E'_1) s_{\lambda^2}(E'_2 - E'_1) \cdots s_{\lambda^p}(E'_p - E'_{p-1})$$

summed over all sequences of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^p)$ with λ^1 k -strict, where $k = n - d_p$. Here Θ_{λ^1} and s_{λ^i} denote theta and Schur polynomials, respectively, and the coefficient $e_{\underline{\lambda}}^w$ is a nonnegative integer which counts the number of p -tuples of leaves of shape $\underline{\lambda}$ in the groves of the transition forest associated to $\mathfrak{d}$ and w .

Let $\text{IG}(n - k, 2n)$ denote the Grassmannian parametrizing isotropic linear subspaces of E of dimension $n - k$. The morphism which sends $E_{\bullet}$ to E_p realizes $\mathfrak{X}(\mathfrak{d})$ as a fiber bundle over $\text{IG}(n - k, 2n)$ with fiber equal to a type A partial flag variety. The mixed nature of the ingredients in formula (2) is in harmony with this fact; however, this simple geometric observation cannot by itself produce an explicit solution to the Giambelli problem.

Our proof of (2) and the corresponding formulas for degeneracy loci is mainly combinatorial. We work with the Schubert polynomials of Billey and Haiman [BH], and more generally with their double versions introduced by Ikeda, Mihalcea, and Naruse [IMN]. These objects have many of the properties of the type A Schubert polynomials, but their connection with the geometry is less clear. This problem was addressed for the single Schubert polynomials in [T2, T3] and in [IMN] for their double counterparts. Using this theory, our geometric results follow readily by combining the Giambelli formulas for isotropic Grassmannians from [BKT2, BKT4] with new *splitting theorems* for these Schubert polynomials, which are analogues of [BKTY, Thm. 4 and Cor. 3] for the other classical Lie types.

A first step towards splitting the Billey-Haiman Schubert polynomials was formulated by Yong [Yo]; however, his result does not suffice for the aforementioned applications to geometry. The point is that the flag of subspaces $E_{\bullet}$ in (1) need not contain a Lagrangian (i.e., a maximal isotropic) subspace. In terms of the Weyl group, this is simply the fact that the first descent position of a signed permutation w need not be at *zero*.

To solve this problem, we introduce the *mixed Stanley functions*, denoted by $J_w(X, Y)$ in type C. The functions J_w are defined as sums of products of type C and type A Stanley symmetric functions in two distinct sets of variables. For each fixed $m \geq 0$, they include among their coefficients the number of reduced words of a signed permutation w such that the last m letters in the word are positive (the letter 0 is used to denote the sign change). When w has no descents smaller than k , where $k \geq m$, the J_w – suitably restricted – are nonnegative integer linear combinations of theta polynomials indexed by k -strict partitions λ . It is the *mixed Stanley coefficients* that appear in this expansion which enter into the splitting and degeneracy locus formulas. Finally, our combinatorial interpretation of the numbers $e_{\underline{\lambda}}^w$ in (2) depends on the *transition equations* of Lascoux and Schützenberger [LS3] and Billey [B]. It would be interesting to have tableau formulas for the $e_{\underline{\lambda}}^w$, analogous to the ones in [BF, BKTY, KMS] for quiver coefficients.

This article is organized so that most of the exposition is in type C; a final section explains the analogous picture in types B and D. We review the Schur, theta, and Schubert polynomials, as well as the Stanley symmetric functions we require in §1. The mixed Stanley functions and their basic properties are studied in §2; this includes applications to enumerating reduced words and product rules for

theta polynomials. Our splitting theorems for Schubert polynomials are proved in §3, and the applications to symplectic degeneracy loci and Giambelli formulas are deduced in §4 and §5, respectively. We conclude with the Schubert splitting results for the orthogonal groups in §6.

I am grateful to my collaborators Anders Buch, Andrew Kresch, and Alexander Yong for their hard work on related papers, especially [BKTY] and [BKT1, BKT2]. I also thank Leonardo Mihalcea for conversations about his joint paper [IMN] on double Schubert polynomials, which played an important role in this project.

1. PRELIMINARIES

1.1. Schur and theta polynomials. An *integer sequence* is a sequence of integers $\alpha = (\alpha_1, \alpha_2, \dots)$ only finitely many of which are non-zero. The largest integer $\ell \geq 0$ such that $\alpha_\ell \neq 0$ is called the *length* of α , denoted $\ell(\alpha)$; we will identify an integer sequence of length ℓ with the vector consisting of its first ℓ terms, and set $|\alpha| = \sum \alpha_i$. An integer sequence λ is a *partition* if $\lambda_i \geq \lambda_{i+1} \geq 0$ for all i . We will represent partitions λ by their Young diagram of boxes, let λ' denote the conjugate (or transpose) of λ , and write $\mu \subset \lambda$ for the containment relation between two Young diagrams. Following Young [Y], given any integer sequence α and natural numbers $i < j$, we define

$$R_{ij}(\alpha) = (\alpha_1, \dots, \alpha_i + 1, \dots, \alpha_j - 1, \dots).$$

A raising operator R is any monomial in these R_{ij} 's. If $(u_1, u_2, \dots)$ is any ordered set of commuting independent variables and α is an integer sequence, we let $u_\alpha = \prod_i u_{\alpha_i}$, with the understanding that $u_0 = 1$ and $u_r = 0$ if $r < 0$. For any raising operator R , let $Ru_\alpha = u_{R\alpha}$.

Let $c = (c_1, c_2, \dots)$ and $d = (d_1, d_2, \dots)$ be two families of commuting variables. Define elements h_r for $r \in \mathbb{Z}$ by the identity of formal power series

$$\sum_{r=-\infty}^{+\infty} h_r t^r = \left(\sum_{i=0}^{\infty} (-1)^i c_i t^i \right)^{-1} \left(\sum_{i=0}^{\infty} (-1)^i d_i t^i \right).$$

Consider the raising operator expression

$$R^0 = \prod_{i < j} (1 - R_{ij})$$

and for any partition λ , define the *Schur polynomial* $s_\lambda(c - d)$ by

$$s_\lambda(c - d) = R^0 h_\lambda = \det(h_{\lambda_i + j - i})_{i,j}.$$

Let $Y = (y_1, y_2, \dots)$ and $Z = (z_1, z_2, \dots)$ be two infinite sets of variables, and define the elementary symmetric functions $e_r(Y)$ by the generating function

$$\prod_{i=1}^{\infty} (1 + y_i t) = \sum_{r=0}^{\infty} e_r(Y) t^r.$$

The *supersymmetric Schur function* $s_\lambda(Y/Z)$ is obtained from $s_\lambda(c - d)$ by setting $c_r = e_r(Y)$ and $d_r = e_r(Z)$ for all $r \geq 1$. The usual Schur S -functions satisfy the identities $s_\lambda(Y) = s_\lambda(Y/Z)|_{Z=0}$ and $s_\lambda(0/Z) = s_\lambda(Y/Z)|_{Y=0} = (-1)^{|\lambda|} s_{\lambda'}(Z)$. In particular, for each integer r , the function $s_r(Y)$ is the complete symmetric function in the variables Y , also denoted $h_r(Y)$.

Fix an integer $k \geq 0$. A partition λ is k -strict if all its parts λ_i greater than k are distinct; λ is called *strict* if it is 0-strict. Any such λ determines a raising operator expression R^λ by the prescription

$$R^\lambda = \prod_{i < j} (1 - R_{ij}) \prod_{\substack{i < j \\ \lambda_i + \lambda_j > 2k + j - i}} (1 + R_{ij})^{-1}$$

where the second product is over all pairs $i < j$ such that $\lambda_i + \lambda_j > 2k + j - i$. Define elements g_r for $r \in \mathbb{Z}$ by the identity

$$\sum_{r=-\infty}^{+\infty} g_r t^r = \left(\sum_{i=0}^{\infty} c_i t^i \right) \left(\sum_{i=0}^{\infty} d_i t^i \right)^{-1}$$

and the *theta polynomial* $\Theta_\lambda(c - d)$ by

$$(3) \quad \Theta_\lambda(c - d) = R^\lambda g_\lambda.$$

If $X = (x_1, x_2, \dots)$ is another infinite set of variables, the formal power series $\vartheta_r(X; Y)$ for $r \in \mathbb{Z}$ are defined by the equation

$$\prod_{i=1}^{\infty} \frac{1 + x_i t}{1 - x_i t} \prod_{j=1}^k (1 + y_j t) = \sum_{r=0}^{\infty} \vartheta_r(X; Y) t^r.$$

Following [BKT2], we then set $\Theta_\lambda(X; Y) = R^\lambda \vartheta_\lambda$. The Θ_λ for λ k -strict form a $\mathbb{Z}$ -basis for the ring $\Gamma^{(k)} = \mathbb{Z}[\vartheta_1, \vartheta_2, \dots]$. The ring $\Gamma = \Gamma^{(0)}$ is the ring of Schur Q -functions (see [M2, III.8] and [Sch]), and in this case $\vartheta_r(X)$ and $\Theta_\lambda(X)$ are denoted by $q_r(X)$ and $Q_\lambda(X)$, respectively.

1.2. The hyperoctahedral group. Let W_n denote the hyperoctahedral group of signed permutations on the set $\{1, \dots, n\}$. We will adopt the notation where a bar is written over an entry with a negative sign. The group W_n is the Weyl group for the root system B_n or C_n , and is generated by the simple transpositions $s_i = (i, i+1)$ for $1 \leq i \leq n-1$ and the sign change $s_0(1) = \bar{1}$. There is a natural embedding $W_n \hookrightarrow W_{n+1}$ defined by adjoining the fixed point $n+1$. The symmetric group S_n is the subgroup of W_n generated by the s_i for $1 \leq i \leq n-1$, and is the Weyl group for the root system A_{n-1} . We let $S_\infty = \cup_n S_n$ and $W_\infty = \cup_n W_n$.

A *reduced word* for $w \in W_n$ is a sequence $a_1 \cdots a_r$ of elements in $\{0, 1, \dots, n-1\}$ such that $w = s_{a_1} \cdots s_{a_r}$ and r is minimal (so equal to the length $\ell(w)$ of w). Given any $u_1, \dots, u_p, w \in W_\infty$, we write $u_1 \cdots u_p = w$ if $\ell(u_1) + \cdots + \ell(u_p) = \ell(w)$ and the product of $u_1, \dots, u_p$ is equal to w . In this case we say that $u_1 \cdots u_p$ is a *reduced factorization* of w . We say that w has *descent* at position $r \geq 0$ if $w_r > w_{r+1}$, where by definition $w_0 = 0$. An element $w \in W_\infty$ is *compatible* with the sequence $\mathbf{a} : a_1 < \cdots < a_p$ of nonnegative integers if all descent positions of w are contained in $\mathbf{a}$. A reduced factorization $u_1 \cdots u_p = w$ is *compatible with $\mathbf{a}$* if $u_j(i) = i$ for all $j > 1$ and $i \leq a_{j-1}$.

We say that a signed permutation $w \in W_\infty$ is *increasing up to k* if it has no descents less than k . This condition is vacuous if $k = 0$, and for positive k it means that $0 < w_1 < w_2 < \cdots < w_k$. An important special case are the k -Grassmannian elements, which by definition satisfy $\ell(ws_i) = \ell(w) + 1$ for all $i \neq k$. There is a natural bijection between k -Grassmannian elements of W_∞ and k -strict

partitions, obtained as follows. If $w \in W_n$ is k -Grassmannian, there exist unique strict partitions u, ζ, v of lengths k, r , and $n - k - r$, respectively, so that

$$w = (u_k, \dots, u_1, \bar{\zeta}_1, \dots, \bar{\zeta}_r, v_{n-k-r}, \dots, v_1).$$

Define μ_i for $1 \leq i \leq k$ by

$$\mu_i = u_i + i - k - 1 + \#\{j \mid \zeta_j > u_i\}.$$

Then w corresponds to the k -strict partition λ such that the lengths of the first k columns of λ are given by $\mu_1, \dots, \mu_k$, and the part of λ in columns $k+1$ and higher is given by ζ . Conversely, for any k -strict λ the corresponding k -Grassmannian element is denoted by w_λ .

1.3. Schubert polynomials and Stanley symmetric functions. Following [FS] and [FK1, FK2], we will use the nilCoxeter algebra $\mathcal{W}_n$ of the hyperoctahedral group W_n to define Schubert polynomials and Stanley symmetric functions in types A and C, respectively. $\mathcal{W}_n$ is the free associative algebra with unity generated by the elements $u_0, u_1, \dots, u_{n-1}$ modulo the relations

$$\begin{aligned} u_i^2 &= 0 & i &\geq 0; \\ u_i u_j &= u_j u_i & |i - j| &\geq 2; \\ u_i u_{i+1} u_i &= u_{i+1} u_i u_{i+1} & i &> 0; \\ u_0 u_1 u_0 u_1 &= u_1 u_0 u_1 u_0. \end{aligned}$$

For any $w \in W_n$, choose a reduced word $a_1 \cdots a_\ell$ for w and define $u_w = u_{a_1} \cdots u_{a_\ell}$. Since the last three relations listed are the Coxeter relations for W_n , it is clear that u_w is well defined, and that the u_w for $w \in W_n$ form a free $\mathbb{Z}$ -basis of $\mathcal{W}_n$. We denote the coefficient of $u_w \in \mathcal{W}_n$ in the expansion of the element $f \in \mathcal{W}_n$ by $\langle f, w \rangle$; thus $f = \sum_{w \in W_n} \langle f, w \rangle u_w$ for all $f \in \mathcal{W}_n$.

Let t be an indeterminate and define

$$\begin{aligned} A_i(t) &= (1 + tu_{n-1})(1 + tu_{n-2}) \cdots (1 + tu_i); \\ \tilde{A}_i(t) &= (1 - tu_i)(1 - tu_{i+1}) \cdots (1 - tu_{n-1}); \\ C(t) &= (1 + tu_{n-1}) \cdots (1 + tu_1)(1 + 2tu_0)(1 + tu_1) \cdots (1 + tu_{n-1}). \end{aligned}$$

According to [FS, Lemma 2.1] and [FK2, Prop. 4.2], for all commuting variables s, t and indices i , the relations $A_i(s)A_i(t) = A_i(t)A_i(s)$ and $C(s)C(t) = C(t)C(s)$ hold. If $C(X) = C(x_1)C(x_2) \cdots$ and $A(Y) = A_1(y_1)A_1(y_2) \cdots$, we deduce that the functions $F_w(X)$ and $G_\varpi(Y)$ defined for $w \in W_n$ and $\varpi \in S_n$ by

$$F_w(X) = \langle C(X), w \rangle \quad \text{and} \quad G_\varpi(Y) = \langle A(Y), \varpi \rangle$$

are symmetric functions in X and Y , respectively. The G_ϖ and F_w are the type A and type C Stanley symmetric functions, introduced in [Sta] and [BH, FK2, L2]. We have that $F_w = F_{w^{-1}}$.

When G_ϖ is expanded in the basis of Schur functions, one obtains a formula

$$(4) \quad G_\varpi(Y) = \sum_{\lambda: |\lambda| = \ell(\varpi)} c_\lambda^\varpi s_\lambda(Y)$$

for some nonnegative integers c_λ^ϖ (see [LS2], [EG]). According to [LS2] (see also [M1, (7.22)]), we have $c_\lambda^\varpi = c_{\lambda'}^{\varpi^{-1}}$. Lascoux and Schützenberger [LS3] gave one of the first combinatorial interpretations for the coefficients c_λ^ϖ , as the number of leaves of shape λ in the *transition tree* $T(\varpi)$ they associated to ϖ . Equation (4) may be used to define the double Stanley symmetric functions $G_\varpi(Y/Z)$.

For any $\varpi \in S_n$, the Schubert polynomial $\mathfrak{S}_\varpi$ of Lascoux and Schützenberger is given by

$$(5) \quad \mathfrak{S}_\varpi(Y) = \langle A_1(y_1)A_2(y_2) \cdots A_{n-1}(y_{n-1}), \varpi \rangle.$$

The definition (5) is equivalent to the one in [LS1], as is shown in [FS]. Now define

$$(6) \quad \mathfrak{C}_w(X; Y, Z) = \left\langle \tilde{A}_{n-1}(z_{n-1}) \cdots \tilde{A}_1(z_1) C(X) A_1(y_1) \cdots A_{n-1}(y_{n-1}), w \right\rangle.$$

If $\mathfrak{C}_w(X; Y) := \mathfrak{C}_w(X; Y, 0)$, then (6) is equivalent to the equation

$$(7) \quad \mathfrak{C}_w(X; Y, Z) = \sum_{uv=w} \mathfrak{S}_{u^{-1}}(-Z) \mathfrak{C}_v(X; Y)$$

summed over all reduced factorizations $uv = w$ with $u \in S_n$. The polynomials $\mathfrak{C}_w$ in (6) were introduced by Ikeda, Mihalcea, and Naruse [IMN]; they are double versions of the type C Billey-Haiman Schubert polynomials [BH]. Their definition differs from (6), but the equivalence of the two follows by combining (7) with [FK2, §7] and [IMN, Cor. 8.10]. One checks that $\mathfrak{S}_\varpi$ and $\mathfrak{C}_w$ are stable under the natural inclusion of W_n in W_{n+1} , and hence well defined for $\varpi \in S_\infty$ and $w \in W_\infty$, respectively. The $\mathfrak{S}_\varpi(Y)$ for $\varpi \in S_\infty$ form a $\mathbb{Z}$ -basis of the polynomial ring $\mathbb{Z}[Y]$, and the $\mathfrak{C}_w(X; Y)$ for $w \in W_\infty$ form a $\mathbb{Z}$ -basis of $\Gamma[Y]$.

If $\varpi \in S_\infty$ is a Grassmannian permutation with a unique descent at r , then the $\mathfrak{S}_\varpi(Y)$ is a Schur polynomial in $(y_1, \dots, y_r)$. In [BKT2, §6], we obtained the analogue of this result for the $\mathfrak{C}_w(X; Y)$: if $w \in W_\infty$ is k -Grassmannian, then $\mathfrak{C}_w(X; Y)$ is a theta polynomial $\Theta_\lambda(X; Y)$, where λ is the k -strict partition associated to w .

1.4. Splitting type A Schubert polynomials. If $r \leq s$ are any two integers, and $P(X, Y, Z)$ is any polynomial or formal power series in the variables x_i , y_j , and z_j , we let $P[r, s]$ denote the power series obtained from $P(X, Y, Z)$ by setting $x_i = 0$ for all i if $0 \notin [r, s]$, $y_j = 0$ if $j \notin [r, s]$, and $z_j = 0$ if $-j \notin [r, s]$. If $0 \in [r, s]$ we set $P^{(r,s)} = P[r, s]$ and $P^{(s)} = P[0, s]$.

If $w \in W_n$ and $v \in S_m$, we define $w \times v \in W_{m+n}$ to be the signed permutation $(w_1, \dots, w_n, v_1 + n, \dots, v_m + n)$. For $\varpi \in S_\infty$ and $1 \leq r \leq s$, equation (5) immediately gives

$$(8) \quad \mathfrak{S}_\varpi[r, s] = \begin{cases} \mathfrak{S}_v(y_r, \dots, y_s) & \text{if } \varpi = 1_{r-1} \times v, \\ 0 & \text{otherwise.} \end{cases}$$

Given any sequence $\mathbf{a} : a_1 < \dots < a_p$ of natural numbers, we furthermore obtain

$$\mathfrak{S}_\varpi(Y) = \sum_{u_1 \cdots u_p = \varpi} \mathfrak{S}_{u_1}[1, a_1] \mathfrak{S}_{u_2}[a_1 + 1, a_2] \cdots \mathfrak{S}_{u_p}[a_{p-1} + 1, a_p]$$

summed over all reduced factorizations $u_1 \cdots u_p = \varpi$.

If ϖ is increasing up to r , then $\mathfrak{S}_\varpi$ is symmetric in $y_1, \dots, y_r$ and we have

$$(9) \quad \mathfrak{S}_\varpi^{(r)} = G_\varpi^{(r)} = \sum_{\lambda : |\lambda| = \ell(\varpi)} c_\lambda^\varpi s_\lambda(y_1, \dots, y_r)$$

with the coefficients c_λ^ϖ as in (4). Suppose now that ϖ is compatible with the sequence $\mathbf{a}$, and set $Y_i = \{y_{a_{i-1}+1}, \dots, y_{a_i}\}$ for each i . From the previous considerations, we deduce that $\mathfrak{S}_\varpi$ satisfies the formula

$$(10) \quad \mathfrak{S}_\varpi(Y) = \sum_{u_1 \cdots u_p = \varpi} G_{u_1}(Y_1) \cdots G_{u_p}(Y_p)$$

summed over all reduced factorizations $u_1 \cdots u_p = \varpi$ compatible with $\mathbf{a}$. Using (4) to refine (10) further gives

$$(11) \quad \mathfrak{S}_\varpi(Y) = \sum_{\underline{\lambda}} c_{\underline{\lambda}}^\varpi s_{\lambda^1}(Y_1) \cdots s_{\lambda^p}(Y_p)$$

summed over all sequences of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^p)$, where

$$c_{\underline{\lambda}}^\varpi = \sum_{u_1 \cdots u_p = \varpi} c_{\lambda^1}^{u_1} \cdots c_{\lambda^p}^{u_p},$$

summed over all reduced factorizations $u_1 \cdots u_p = \varpi$ compatible with $\mathbf{a}$. More general versions of (10), (11) for universal Schubert polynomials [Fu4] and quiver polynomials [BF] are established in [BKTY, KMS].

Following [BKTY, Cor. 3], a result of Fomin and Greene [FG] for the coefficients c_λ^ϖ in (4) implies an alternative formula for the $c_{\underline{\lambda}}^\varpi$. The *column word* $\text{col}(T)$ of a semistandard Young tableau T is obtained by reading the entries in the columns of T from bottom to top and left to right. For any $\varpi \in S_\infty$ compatible with $\mathbf{a}$, the coefficient $c_{\underline{\lambda}}^\varpi$ equals the number of sequences of semistandard tableaux $(T_1, \dots, T_p)$ such that $\text{col}(T_1) \cdots \text{col}(T_p)$ is a reduced word for ϖ , the entries of T_i are strictly greater than a_{i-1} and the shape of T_i is conjugate to λ^i for each i .

2. TRANSITION FOR MIXED STANLEY FUNCTIONS

2.1. Mixed Stanley functions.

Definition 1. Given $w \in W_\infty$, the (right) *mixed Stanley function* $J_w(X; Y)$ is defined by the equation

$$J_w(X; Y) = \langle C(X)A(Y), w \rangle = \sum_{uv=w} F_u(X)G_v(Y)$$

summed over all reduced factorizations $uv = w$ with $v \in S_\infty$.

Definition 1 can be easily restated in terms of reduced decompositions and admissible sequences, along the lines of [BH, Eq. (3.2)]. One has a dual notion of a left mixed Stanley function $J'_w(X; Y) = \sum_{uv=w} G_{u^{-1}}(Y)F_v(X)$, summed over all reduced factorizations $uv = w$ with $u \in S_\infty$. This is equivalent to the right version since clearly $J'_w(X; Y) = J_{w^{-1}}(X, Y)$.

Recall that $J_w^{(k)} = J_w[0, k]$. If $w \in W_\infty$ is increasing up to k , we claim that $\mathfrak{C}_w^{(k)} = J_w^{(k)}$. Indeed, observe that if $w = uv$ is a reduced factorization, then $\ell(vs_i) = \ell(v) + 1$ for all $i < k$, i.e., v is also increasing up to k . It follows that

$$(12) \quad \mathfrak{C}_w^{(k)} = \sum_{uv=w, v \in S_\infty} F_u(X)\mathfrak{S}_v^{(k)} = \sum_{uv=w, v \in S_\infty} F_u(X)G_v^{(k)} = J_w^{(k)},$$

as claimed. In particular, if $w = w_\lambda$ is k -Grassmannian, then

$$(13) \quad J_{w_\lambda}^{(k)}(X; Y) = \Theta_\lambda(X; Y).$$

Example 1. If λ is a k -strict partition, then

$$(14) \quad J_{w_\lambda}(X; Y) = \sum_{\mu \subset \lambda} F_{\lambda/\mu}(X) s_{\mu'}(Y)$$

summed over all k -strict partitions $\mu \subset \lambda$. The function $F_{\lambda/\mu}(X)$ in (14) is the skew F -function from [T2, §5], which, when non-zero, is equal to $F_{w_\lambda w_\mu^{-1}}(X)$.

2.2. Transition equations. For positive integers $i < j$ we define reflections $t_{ij} \in S_\infty$ and $\bar{t}_{ij}, \bar{t}_{ii} \in W_\infty$ by their right actions

$$\begin{aligned} (\dots, w_i, \dots, w_j, \dots) t_{ij} &= (\dots, w_j, \dots, w_i, \dots), \\ (\dots, w_i, \dots, w_j, \dots) \bar{t}_{ij} &= (\dots, \bar{w}_j, \dots, \bar{w}_i, \dots), \quad \text{and} \\ (\dots, w_i, \dots) \bar{t}_{ii} &= (\dots, \bar{w}_i, \dots). \end{aligned}$$

We let $\bar{t}_{ji} = \bar{t}_{ij}$. According to [B, Thms. 4, 5], the type C Schubert polynomials $\mathfrak{C}_w = \mathfrak{C}_w(X; Y)$ satisfy the recursion formula

$$(15) \quad \mathfrak{C}_w = y_r \mathfrak{C}_{wt_{rs}} + \sum_{\substack{1 \leq i < r \\ \ell(wt_{rs}t_{ir}) = \ell(w)}} \mathfrak{C}_{wt_{rs}t_{ir}} + \sum_{\substack{i \geq 1 \\ \ell(wt_{rs}\bar{t}_{ir}) = \ell(w)}} \mathfrak{C}_{wt_{rs}\bar{t}_{ir}},$$

where r is the last positive descent of w and s is maximal such that $w_s < w_r$. If the last descent r of w satisfies $r > k$, we deduce from (12) and (15) that

$$(16) \quad J_w^{(k)} = \sum_{\substack{1 \leq i < r \\ \ell(wt_{rs}t_{ir}) = \ell(w)}} J_{wt_{rs}t_{ir}}^{(k)} + \sum_{\substack{i \geq 1 \\ \ell(wt_{rs}\bar{t}_{ir}) = \ell(w)}} J_{wt_{rs}\bar{t}_{ir}}^{(k)}.$$

For any $w \in W_\infty$ which is increasing up to k , we construct a rooted tree $T^k(w)$ with nodes given by elements of W_∞ and root w as follows. Let r be the last descent of w . If $w = 1$ or $r = k$, then set $T^k(w) = \{w\}$. Otherwise, let $s = \max\{i > r \mid w_i < w_r\}$ and $\Phi(w) = \Phi_1(w) \cup \Phi_2(w)$, where

$$\begin{aligned} \Phi_1(w) &= \{wt_{rs}t_{ir} \mid 1 \leq i < r \text{ and } \ell(wt_{rs}t_{ir}) = \ell(w)\}, \\ \Phi_2(w) &= \{wt_{rs}\bar{t}_{ir} \mid i \geq 1 \text{ and } \ell(wt_{rs}\bar{t}_{ir}) = \ell(w)\}. \end{aligned}$$

To recursively define $T^k(w)$, we join w by an edge to each $v \in \Phi(w)$, and attach to each $v \in \Phi(w)$ its tree $T^k(v)$. We call $T^k(w)$ the k -transition tree of w .

Lemma 1. *The k -transition tree $T^k(w)$ is finite. All the nodes of $T^k(w)$ are increasing up to k , and the leaves of $T^k(w)$ are k -Grassmannian elements.*

Proof. If w is k -Grassmannian then clearly $T^k(w) = w$. We will show that if w is increasing up to k and not k -Grassmannian, then $\Phi(w) \neq \emptyset$ and all elements of $\Phi(w)$ are increasing up to k . Let $r > k$ be the last descent of w , s be maximal such that $w_s < w_r$, and $v = wt_{rs}$. If $\Phi_1(w) = \emptyset$, then according to Monk's criterion [Mo] (see also [B, Lemma 1]), we must have $w_i > w_s$ for all $i < r$. If $w_s > 0$ then $v\bar{t}_{rr} \in \Phi_2(w)$. If $w_s < 0$, let $i > 0$ be minimal such that $w_i > -w_s$. Then one can easily check using [B, Lemma 2] that $v\bar{t}_{ir} \in \Phi_2(w)$. This proves that $\Phi(w)$ is nonempty.

To show that all elements of $\Phi(w)$ are increasing up to k , we may assume that $k > 0$. If there exists an $i \leq k$ such that $\ell(vt_{ir}) = \ell(w)$, then $w_s > w_i$ and for all $j \in (i, r)$, $w_j \notin (w_i, w_s)$. It follows that vt_{ir} is increasing up to k . If $\ell(v\bar{t}_{ir}) = \ell(w)$ for some $i \leq k$, then we must have $w_s < 0$. Moreover, by [B, Lemma 3], the descent

$$(18) \quad F_w(X) = \sum_{\lambda: |\lambda|=\ell(w)} e_\lambda^w Q_\lambda(X)$$

summed over strict partitions λ . A transition based formula for the constants e_λ^w in equation (18) was proved by Billey [B]. There are several alternative combinatorial descriptions of these numbers in this case, which include a formula in terms of *Kraskiewicz tableaux* [Kr, L2] of shape λ . It would be interesting to have an analogous tableau formula for the e_λ^w in the general case where $k > 0$. Another natural question is whether the mixed Stanley coefficients can be used to obtain a Littlewood-Richardson type rule for theta polynomials; some positive results in this direction are explained in §2.4.

Example 3. Consider the *double mixed Stanley function*

$$J_w(X; Y/Z) = \sum_{\varpi uv=w} G_{\varpi^{-1}}(-Z) F_u(X) G_v(Y)$$

summed over all reduced factorizations $\varpi uv = w$ with $\varpi, v \in S_\infty$. Fix integers $j, k \geq 0$, and suppose that $w \in W_\infty$ is increasing up to k and w^{-1} is increasing up to j . Then $\mathfrak{C}_w^{(-j,k)}(X; Y, Z) = J_w^{(-j,k)}(X; Y/Z)$. However, the analogue of Theorem 1 fails, at least for fixed k . For an example with $j = k = 1$, one checks that $\mathfrak{C}_{231}^{(-1,1)}$ cannot be written as an integral linear combination of $\mathfrak{C}_{2\bar{1}3}^{(-1,1)}$ and $\mathfrak{C}_{312}^{(-1,1)}$, while $2\bar{1}3$ and 312 are the only 1-Grassmannian elements of length two in W_∞ .

2.3. Reduced words and k -bitableaux. The type A Stanley symmetric functions G_ϖ were used in [Sta] and to express the number of reduced words of a permutation ϖ in terms of the numbers f^λ of standard tableaux of shape λ , for the partitions λ which appear in equation (4). Similarly, the type C Stanley symmetric functions F_w can be used to compute the number of reduced words for an element $w \in W_\infty$, as shown in [H, Kr]. We proceed to give an analogue of these results for the mixed Stanley functions J_w .

We first recall some definitions from [T2, §5]. Let $\mathbf{P}$ denote the ordered alphabet $\{1' < 2' < \dots < k' < 1 < 2 < \dots\}$. The symbols $1', \dots, k'$ are called *marked*, while the rest are *unmarked*. Let λ be a k -strict partition. A k -bitableau U of shape λ is a filling of the boxes in λ with elements of $\mathbf{P}$ which is weakly increasing along each row and down each column, such that (i) the marked entries are strictly increasing along rows and weakly increasing down columns, and (ii) the unmarked entries form a k -tableau T . Each k -bitableau U has an associated multiplicity $r(U)$; see loc. cit. for definitions and more details. Let $(xy)^U = \prod_i x_i^{m_i} \prod_j y_j^{n_j}$, where m_i (respectively n_j) denotes the number of times that i (respectively j') appears in U . According to [T2, Thm. 5], we have

$$(19) \quad \Theta_\lambda(X; Y) = \sum_U 2^{r(U)} (xy)^U$$

summed over all k -bitableaux U of shape λ .

If a k -bitableau U contains exactly m marked entries, we say that U is of *type* m . U is called *standard* if the entries $1', \dots, m', 1, \dots, n$ each appear once in U for some n ; in this case we have $r(U) = n$.

Definition 3. Let m be a nonnegative integer with $m \leq k$. For any k -strict partition λ , define h_m^λ to be the number of standard k -bitableaux of shape λ and type m . We say that a reduced word for $w \in W_\infty$ has *type* m if the last m letters of the word are positive.

Following [T2, Thm. 6 and Ex. 9], for any two k -strict partitions λ, μ with $\mu \subset \lambda$ and $\mu_1 \leq k$, the number $g^{\lambda/\mu}$ of standard k -tableaux of shape λ/μ is equal to the number of reduced words for $w_\lambda w_\mu^{-1}$. Furthermore, using e.g. [T2, Prop. 5], one can construct a bijection between the set of reduced words of type m for w_λ and the set of k -bitableaux of shape λ and type m , for any k -strict partition λ . It follows that for any $m \leq k$, we have

$$h_m^\lambda = \sum_{\mu \subset \lambda, |\mu|=m} f^\mu g^{\lambda/\mu}$$

summed over all partitions $\mu \subset \lambda$ with $|\mu| = m$. The $k = 0$ case of the next result is due to Haiman [H] and Kraśkiewicz [Kr].

Proposition 1. *Let $w \in W_\infty$ be increasing up to k and let m be an integer with $0 \leq m \leq \min(k, \ell(w))$. Then the number of reduced words of type m for w is equal to $\sum_\lambda e_\lambda^w h_m^\lambda$, where the sum is over all k -strict partitions λ .*

Proof. It is clear that the number of reduced words of type m for w equals 2^{-n} times the coefficient of $x_1 \cdots x_n y_1 \cdots y_m$ in J_w , where $n = \ell(w) - m$. On the other hand, this coefficient is also equal to $2^n \sum_\lambda e_\lambda^w h_m^\lambda$, by (17) and (19). $\square$

It would be interesting to find a bijective proof of Proposition 1.

2.4. Multiplication rules. We show here how Theorem 1 and Lemma 2 lead to Littlewood-Richardson type rules for the product of two theta polynomials. In the case of $k = 0$, we observe that the transition equations in [B] give a combinatorial rule for the structure constants in the product of *any* two Schur Q -functions. This answers a question of Manivel in the affirmative (compare with [BL, §4]).

We will actually work with the Schur P -functions, which are defined by the equation $P_\lambda = 2^{-\ell(\lambda)} Q_\lambda$, for any strict partition λ . Given two strict partitions μ and ν , there are nonnegative integers $f_{\mu\nu}^\lambda$ such that

$$P_\mu P_\nu = \sum_\lambda f_{\mu\nu}^\lambda P_\lambda.$$

The $f_{\mu\nu}^\lambda$ agree with the Schubert structure constants on maximal orthogonal Grassmannians $\text{OG}(n, 2n+1)$, when n is sufficiently large. The same numbers also appear in the expansion of skew Schur Q -functions

$$(20) \quad Q_{\lambda/\mu} = \sum_\nu f_{\mu\nu}^\lambda Q_\nu$$

in the Q -basis. Combinatorial rules for the $f_{\mu\nu}^\lambda$ may be found in [Sa, W, St1, Sh].

The skew Schur Q -functions are known to be equal to certain type C Stanley symmetric functions (see [FK2, Thm. 8.2] and [St2, Cor. 6.6]). In fact, we have $Q_{\lambda/\mu} = F_{w_\lambda w_\mu^{-1}}$, where w_λ and w_μ are the 0-Grassmannian elements associated to λ and μ (this is a special case of [T2, Thm. 6]). We therefore deduce the following result from (18) and (20).

Corollary 1. *The coefficient $f_{\mu\nu}^\lambda$ is equal to the number of leaves of $T^0(w_\lambda w_\mu^{-1})$ of shape ν , if $\mu \subset \lambda$, and is equal to zero, otherwise.*

Now let k be any nonnegative integer. The following result is an analogue of [BL, Lemma 16] for the functions J_w .

Lemma 2. For $w \in W_n$ and $v \in S_\infty$, we have

$$J_w J_v = J_{w \times v}.$$

Proof. Since $a_1 \cdots a_\ell$ is a reduced word for $v \in S_m$ if and only if $(a_1 + n) \cdots (a_\ell + n)$ is a reduced word for $1_n \times v$, we see that $G_{1_n \times v} = G_v$, $F_{1_n \times v} = F_v$, and $J_v = J_{1_n \times v}$ for each $n \geq 1$. The reduced words for $w \times v$ are all obtained by intertwining a reduced word for w with a reduced word for $1_n \times v$. Moreover, given any reduced factorization $ab = w \times v$, with $b \in S_\infty$, we have $a = a_1 \times a_2$ and $b = b_1 \times b_2$ where $a_1, b_1 \in W_n$ and $a_2, b_1, b_2 \in S_\infty$. We deduce that

$$\begin{aligned} J_{w \times v}(X; Y) &= \sum_{(a_1 \times a_2)(b_1 \times b_2) = w \times v} F_{a_1 \times a_2}(X) G_{b_1 \times b_2}(Y) \\ &= \sum_{(a_1 \times a_2)(b_1 \times b_2) = w \times v} F_{a_1}(X) F_{1_n \times a_2}(X) G_{b_1}(Y) G_{1_n \times b_2}(Y) \\ &= \sum_{a_1 b_1 = w, a_2 b_2 = v} F_{a_1}(X) G_{b_1}(Y) F_{a_2}(X) G_{b_2}(Y) \\ &= J_w(X; Y) J_v(X; Y). \end{aligned} \quad \square$$

We obtain a combinatorial rule for multiplying two theta polynomials, when one of the factors is indexed by a ‘small’ partition (compare with [BL, Cor. 17]).

Corollary 2. Let μ and ν be k -strict partitions with $\nu_i \leq k$ for all i , and consider the product expansion

$$\Theta_\mu \Theta_\nu = \sum_{\lambda} \varphi_{\mu\nu}^\lambda \Theta_\lambda$$

summed over k -strict partitions λ . Then $\varphi_{\mu\nu}^\lambda$ is equal to the number of leaves of $T^k(w_\mu \times w_\nu)$ of shape λ .

Proof. Observe that $w_\nu \in S_\infty$ if and only if $\nu_i \leq k$ for all i . Lemma 2 therefore applies and gives the equation

$$\Theta_\mu \Theta_\nu = J_{w_\mu}^{(k)} J_{w_\nu}^{(k)} = J_{w_\mu \times w_\nu}^{(k)}.$$

The result now follows from Theorem 1. Recall from [BKT2, §5.4] that in this situation we have $\Theta_\nu = R^0 \vartheta_\nu = \det(\vartheta_{\nu_i + j - i})_{i,j}$. $\square$

Example 4. For the 1-strict partitions $\mu = (2, 1)$ and $\nu = 1$ we have $w_\mu \times w_\nu = 3\bar{1}254 \in W_5$. Example 2 therefore gives $\Theta_{(2,1)} \Theta_1 = \Theta_{(2,1,1)} + 2 \Theta_{(3,1)} + \Theta_4$.

3. SPLITTING TYPE C SCHUBERT POLYNOMIALS

In this section we give splitting theorems for the single and double type C Schubert polynomials $\mathfrak{C}_w$. For any $k \geq 0$, let $Y_{>k} = (y_{k+1}, y_{k+2}, \dots)$. The following proposition generalizes the $k = 0$ case from [BH, Thm. 3].

Proposition 2. If $w \in W_\infty$ is increasing up to k , then

$$(21) \quad \mathfrak{C}_w(X; Y) = \sum_{u(1_k \times v) = w} J_u^{(k)}(X; Y) \mathfrak{S}_v(Y_{>k})$$

where the sum is over all reduced factorizations $u(1_k \times v) = w$ with $v \in S_\infty$. The Schubert polynomials $\mathfrak{C}_w(X; Y)$ for $w \in W_\infty$ increasing up to k form a $\mathbb{Z}$ -basis for the ring $\Gamma^{(k)}[Y_{>k}] = \Gamma^{(k)}[y_{k+1}, y_{k+2}, \dots]$.

Proof. From the definition (6) we deduce that for any $w \in W_n$,

$$\mathfrak{C}_w(X; Y) = \sum_{uv=w, v \in S_\infty} \mathfrak{C}_u[0, k] \mathfrak{S}_v[k+1, n].$$

According to (8), the polynomial $\mathfrak{S}_v[k+1, n]$ is non-zero only if $v = 1_k \times \varpi$ for some $\varpi \in S_\infty$, in which case $\mathfrak{S}_v[k+1, n] = \mathfrak{S}_\varpi(y_{k+1}, \dots, y_n)$. For all such v , we furthermore note that the element $u = wv^{-1}$ is increasing up to k , and hence $\mathfrak{C}_u[0, k] = J_u[0, k]$ by Theorem 1. This proves equation (21).

Set $\mathfrak{C}_w = \mathfrak{C}_w(X; Y)$ and for each $i \geq 0$, let ∂_i denote the divided difference operator from [BH, §2]. Recall that $\partial_i \mathfrak{C}_w = \mathfrak{C}_{ws_i}$, if $w_i > w_{i+1}$, and $\partial_i \mathfrak{C}_w = 0$, otherwise. Since the $\Theta_\lambda = \mathfrak{C}_{w_\lambda}$ for k -strict partitions λ form a $\mathbb{Z}$ -basis of $\Gamma^{(k)}$, we deduce that $\partial_i f = 0$ for all $i < k$ and f in the ring $\Gamma^{(k)}[Y_{>k}]$.

Equations (17) and (21) imply that the $\mathfrak{C}_w$ for w increasing up to k are contained in $\Gamma^{(k)}[Y_{>k}]$. Moreover, the $\mathfrak{C}_w$ for $w \in W_\infty$ are known to be a $\mathbb{Z}$ -basis of $\Gamma[Y]$ from [BH, Thm. 3]. Given any $f \in \Gamma^{(k)}[Y_{>k}]$, we therefore have $f = \sum_{w \in W_\infty} a_w \mathfrak{C}_w$ for some $a_w \in \mathbb{Z}$. Since $\partial_i f = 0$ for all $i < k$, we deduce that only terms $\mathfrak{C}_w$ with w increasing up to k appear in the sum, completing the proof. $\square$

Fix a sequence $\mathbf{a} : a_1 < \dots < a_p$ of nonnegative integers and define $Y_i = \{y_{a_{i-1}+1}, \dots, y_{a_i}\}$ for each i ; in particular $Y_1 = \emptyset$ if $a_1 = 0$. Given any sequence of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^p)$ with λ^1 a_1 -strict, we define the nonnegative integer

$$(22) \quad e_{\underline{\lambda}}^w = \sum_{u_1 \cdots u_p = w} e_{\lambda^1}^{u_1} c_{\lambda^2}^{u_2} \cdots c_{\lambda^p}^{u_p},$$

where the sum is over reduced factorizations $u_1 \cdots u_p = w$ compatible with $\mathbf{a}$ such that $u_2, \dots, u_p \in S_\infty$, and the integers $c_{\lambda^i}^{u_i}$ and $e_{\lambda^1}^{u_1}$ are as in (4) and (17), respectively.

The number $e_{\underline{\lambda}}^w$ can be described in a more picturesque way as follows. Given a permutation $\varpi \in S_\infty$, let $T(\varpi)$ denote the Lascoux-Schützenberger transition tree associated to ϖ in [LS3, §4]. For any reduced factorization $u_1 \cdots u_p = w$ compatible with $\mathbf{a}$ such that $u_2, \dots, u_p \in S_\infty$, the p -tuple of trees $(T^{a_1}(u_1), T(u_2), \dots, T(u_p))$ is called a *grove*. The collection of all such groves forms the $\mathbf{a}$ -transition forest associated to w . The integer $e_{\underline{\lambda}}^w$ is then equal to the number of p -tuples of leaves of shape $\underline{\lambda}$ in the groves of the $\mathbf{a}$ -transition forest associated to w .

Theorem 2. *Suppose that $w \in W_\infty$ is compatible with the sequence $\mathbf{a}$. Then we have*

$$(23) \quad \mathfrak{C}_w(X; Y) = \sum_{u_1 \cdots u_p = w} J_{u_1}(X; Y_1) G_{u_2}(Y_2) \cdots G_{u_p}(Y_p)$$

summed over all reduced factorizations $u_1 \cdots u_p = w$ compatible with $\mathbf{a}$ such that $u_2, \dots, u_p \in S_\infty$. Furthermore, we have

$$(24) \quad \mathfrak{C}_w(X; Y) = \sum_{\underline{\lambda}} e_{\underline{\lambda}}^w \Theta_{\lambda^1}(X; Y_1) s_{\lambda^2}(Y_2) \cdots s_{\lambda^p}(Y_p)$$

summed over all sequences of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^p)$ with λ^1 a_1 -strict.

Proof. Equation (23) follows from (21) and the type A Schubert splitting formula (10). Moreover, (24) is obtained from (23) by using equations (8), (9), and (17). $\square$

Let $\underline{\lambda}^- = (\lambda^2, \dots, \lambda^p)$. An equivalent expression for the coefficients $e_{\underline{\lambda}}^w$ is

$$(25) \quad e_{\underline{\lambda}}^w = \sum_{uv=w} e_{\lambda^1}^u c_{\underline{\lambda}^-}^v$$

summed over all reduced factorizations $uv = w$ with $v \in S_\infty$, where $c_{\underline{\lambda}^-}^v$ equals the number of sequences of tableaux $(T_2, \dots, T_p)$ such that $\text{col}(T_2) \cdots \text{col}(T_p)$ is a reduced word for v , the entries of T_i are strictly greater than a_{i-1} and the shape of T_i is conjugate to λ^i for each $i \geq 2$. With this interpretation of the numbers $e_{\underline{\lambda}}^w$, the $k = 0$ case of formula (24) is contained in [Yo, §5].

Fix a second sequence $\mathbf{b} : 0 = b_1 < \dots < b_q$ of nonnegative integers and set $Z_j = \{z_{b_{j-1}+1}, \dots, z_{b_j}\}$ for each j . A reduced factorization $u_1 \cdots u_{p+q-1} = w$ is *compatible with $\mathbf{a}, \mathbf{b}$* if $u_j(i) = i$ whenever $j < q$ and $i \leq b_{q-j}$ or whenever $j > q$ and $i \leq a_{j-q}$. Given any sequence of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^{p+q-1})$ with λ^q a_1 -strict, we define

$$(26) \quad f_{\underline{\lambda}}^w = \sum_{u_1 \cdots u_{p+q-1} = w} c_{\lambda^1}^{u_1} \cdots c_{\lambda^{q-1}}^{u_{q-1}} e_{\lambda^q}^{u_q} c_{\lambda^{q+1}}^{u_{q+1}} \cdots c_{\lambda^{p+q-1}}^{u_{p+q-1}},$$

where the sum is over reduced factorizations $u_1 \cdots u_{p+q-1} = w$ compatible with $\mathbf{a}, \mathbf{b}$ such that $u_i \in S_\infty$ for all $i \neq q$. Using (7) and (11), one can easily obtain an equivalent formula for the $f_{\underline{\lambda}}^w$ along the lines of (25).

Corollary 3. *Suppose that w and w^{-1} are compatible with the sequences $\mathbf{a}$ and $\mathbf{b}$, respectively. Then $\mathfrak{C}_w(X; Y, Z)$ is equal to*

$$\sum_{u_1 \cdots u_{p+q-1} = w} G_{u_1}(0/Z_q) \cdots G_{u_{q-1}}(0/Z_2) J_{u_q}(X; Y_1) G_{u_{q+1}}(Y_2) \cdots G_{u_{p+q-1}}(Y_p)$$

summed over all reduced factorizations $u_1 \cdots u_{p+q-1} = w$ compatible with $\mathbf{a}, \mathbf{b}$ such that $u_i \in S_\infty$ for all $i \neq q$. Furthermore, we have

$$(27) \quad \mathfrak{C}_w(X; Y, Z) = \sum_{\underline{\lambda}} f_{\underline{\lambda}}^w s_{\lambda^1}(0/Z_q) \cdots \Theta_{\lambda^q}(X; Y_1) \cdots s_{\lambda^{p+q-1}}(Y_p)$$

summed over all sequences of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^{p+q-1})$ with λ^q a_1 -strict.

Proof. The result follows immediately from equations (7), (10), and Theorem 2. $\square$

According to [IMN, Thm. 8.1], the Schubert polynomials $\mathfrak{C}_w(X; Y, Z)$ enjoy the following symmetry property:

$$(28) \quad \mathfrak{C}_w(X; Y, Z) = \mathfrak{C}_{w^{-1}}(X; -Z, -Y).$$

This also follows immediately from equation (6). By applying Corollary 3 to the right hand side of (28), we obtain dual versions of these splitting equations.

4. SYMPLECTIC DEGENERACY LOCI

4.1. Isotropic partial flag bundles. Let $E \rightarrow B$ be a vector bundle of rank $2n$ on an algebraic variety B . Assume that E is a *symplectic* bundle, i.e. E is equipped with an everywhere nondegenerate skew-symmetric form $E \otimes E \rightarrow \mathbb{C}$. A subbundle V of E is *isotropic* if the form vanishes when restricted to V ; the ranks of isotropic subbundles of E range from 0 to n . Fix a sequence $\mathbf{a} : a_1 < \dots < a_p$ of nonnegative integers with $a_p < n$, and set $a_{p+1} = n$ for convenience. We introduce the *isotropic*

partial flag bundle $F^a(E)$ with its projection map $\rho : F^a(E) \rightarrow \mathfrak{X}$. The variety $F^a(E)$ parametrizes partial flags

$$(29) \quad E_\bullet : 0 = E_0 \subset E_1 \subset \cdots \subset E_p \subset E$$

with $\text{rank } E_i = n - a_{p+1-i}$ and E_p isotropic. Here we have identified E with its pullback under the map ρ , and also let (29) denote the tautological partial flag of vector bundles over $F^a(E)$.

There is a group monomorphism $\phi : W_n \hookrightarrow S_{2n}$ with image

$$\phi(W_n) = \{ \varpi \in S_{2n} \mid \varpi(i) + \varpi(2n+1-i) = 2n+1, \text{ for all } i \}.$$

The map ϕ is determined by setting, for each $w = (w_1, \dots, w_n) \in W_n$ and $1 \leq i \leq n$,

$$\phi(w)(i) = \begin{cases} n+1-w_{n+1-i} & \text{if } w_{n+1-i} \text{ is unbarred,} \\ n+\overline{w}_{n+1-i} & \text{otherwise.} \end{cases}$$

Let W^a be the set of signed permutations $w \in W_n$ whose descent positions are listed among the integers $a_1, \dots, a_p$. These elements are the minimal length coset representatives in W_n/W_a , where W_a denotes the subgroup of W_n generated by the simple reflections s_i for $i \notin \{a_1, \dots, a_p\}$.

Fix a flag $0 = F_0 \subset F_1 \subset \cdots \subset F_n \subset E$ of subbundles of E with $\text{rank } F_i = i$ for each i and F_n isotropic. We extend any such flag to a complete flag $F_\bullet$ in E by letting $F_{n+i} = F_{n-i}^\perp$ for $1 \leq i \leq n$; we call the completed flag a *complete isotropic flag*. For every $w \in W^a$ and complete isotropic flag $F_\bullet$ of subbundles of E , we define the *universal Schubert variety* $\mathfrak{X}_w \subset F^a(E)$ as the locus of $a \in F^a(E)$ such that

$$(30) \quad \dim(E_r(a) \cap F_s(a)) \geq \# \{ i \leq r \mid \phi(w)(i) > 2n-s \} \quad \forall r, s.$$

The variety $\mathfrak{X}_w$ is an irreducible subvariety of $F^a(E)$ of codimension $\ell(w)$, and may be regarded as a universal degeneracy locus. Formulas for the classes $[\mathfrak{X}_w]$ in the cohomology or Chow ring of $F^a(E)$ pull back to identities for corresponding loci whenever one has a symplectic vector bundle V and two flags of isotropic subbundles of V , following [Fu3]. Moreover, they are equivalent to formulas which represent the Schubert classes in the T -equivariant cohomology ring of isotropic partial flag varieties, as observed e.g. in [Gra].

4.2. Full flag bundles and the geometrization map. Consider the full flag bundle $F(E) = F^{(0,1,2,\dots,n-1)}(E)$ parametrizing complete isotropic flags of subbundles $E_\bullet$ in E . Let $X = (x_1, \dots, x_n)$ and $Y = (y_1, \dots, y_n)$. According to [Fu2, §3], the cohomology (or Chow) ring $H^*(F(E), \mathbb{Z})$ is presented as a quotient

$$(31) \quad H^*(F(E)) \cong H^*(B)[X, Y]/J_n,$$

where J_n denotes the ideal generated by the differences $e_i(x_1^2, \dots, x_n^2) - e_i(y_1^2, \dots, y_n^2)$ for $1 \leq i \leq n$. The inverse of the isomorphism (31) sends the class of x_i to $-c_1(E_{n+1-i}/E_{n-i})$ and of y_i to $-c_1(F_{n+1-i}/F_{n-i})$ for each i with $1 \leq i \leq n$.

The ring $H^*(F(E))$ may be used to study the cohomology of any isotropic partial flag bundle $F^a(E)$, because the natural surjection $F(E) \rightarrow F^a(E)$ induces an injective ring homomorphism $\iota : H^*(F^a(E)) \rightarrow H^*(F(E))$. The tautological vector bundles E_i, F_j , the universal Schubert varieties, and their cohomology classes on $F^a(E)$ pull back under ι to the homonymous objects over $F(E)$.

The Schubert varieties $\mathfrak{X}_w$ on $F(E)$ are indexed by w in the full Weyl group W_n . Furthermore, the type C double Schubert polynomials $\mathfrak{C}_w(X; Y, Z)$ represent

their cohomology classes $[\mathfrak{X}_w]$ in the presentation (31), but only after a certain change of variables. Ikeda, Mihalcea, and Naruse [IMN, §10] provide a different way to connect these Schubert polynomials to geometry, which we will adapt to our current setup. For a closely related but distinct approach in the case of single Schubert polynomials, see [T1].

The key tool is the following ring homomorphism from [IMN], which we call the *geometrization map*:

$$\pi_n : \Gamma[Y, Z] \rightarrow H^*(B)[X, Y]/J_n.$$

The homomorphism π_n is determined by setting

$$\pi_n(q_r(X)) = \sum_{i=0}^r e_i(X) h_{r-i}(Y) \quad \text{for all } r,$$

$$\pi_n(y_i) = -x_i \quad \text{for } 1 \leq i \leq n, \quad \pi_n(y_i) = 0 \quad \text{for } i > n, \quad \pi_n(z_j) = y_j \quad \text{for each } j.$$

It follows from [IMN, Prop. 7.7] that for $w \in W_n$, the geometrization map π_n maps $\mathfrak{C}_w(X; Y, Z)$ to a polynomial which represents the universal Schubert class $[\mathfrak{X}_w]$ in the presentation (31). Furthermore, we have $\pi_n(\mathfrak{C}_w) \in J_n$ for $w \in W_\infty \setminus W_n$.

If V and V' are two vector bundles with total Chern classes $c(V)$ and $c(V')$, respectively, we denote $s_\lambda(c(V) - c(V'))$ by $s_\lambda(V - V')$. We similarly denote the class $\Theta_\lambda(c(V) - c(V'))$ by $\Theta_\lambda(V - V')$. To state our main geometric result, let $Q_1 = E/E_p, Q_2 = E_p/E_{p-1}, \dots, Q_p = E_2/E_1$. Consider a sequence $\mathfrak{b} : 0 = b_1 < \dots < b_q$ with $b_q < n$, and set $\hat{Q}_2 = F_n/F_{n-b_2}, \dots, \hat{Q}_q = F_{n-b_{q-1}}/F_{n-b_q}$.

Theorem 3. *Suppose that $w \in W^a$ and that w^{-1} is compatible with $\mathfrak{b}$. Then we have*

$$\begin{aligned} [\mathfrak{X}_w] &= \sum_{\underline{\lambda}} f_{\underline{\lambda}}^w s_{(\lambda^1)'}(\hat{Q}_q) \cdots s_{(\lambda^{q-1})'}(\hat{Q}_2) \Theta_{\lambda^q}(Q_1 - F_n) s_{\lambda^{q+1}}(Q_2) \cdots s_{\lambda^{p+q-1}}(Q_p) \\ &= \sum_{\underline{\lambda}} f_{\underline{\lambda}}^w s_{\lambda^1}(F_{n+b_{q-1}} - F_{n+b_q}) \cdots \Theta_{\lambda^q}(E - E_p - F_n) \cdots s_{\lambda^{p+q-1}}(E_2 - E_1), \end{aligned}$$

where the sum is over all sequences of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^{p+q-1})$ with λ^q a_1 -strict, and the coefficients $f_{\underline{\lambda}}^w$ are given by (26).

Proof. The variables x_i for $1 \leq i \leq n$ give the Chern roots of the various vector bundles over $F^a(E)$. In particular the Chern roots of Q_1 are $x_1, \dots, x_n, -x_1, \dots, -x_{a_1}$, while those of Q_r for $r \geq 2$ are $-x_{a_{r-1}+1}, \dots, -x_{a_r}$. Similarly the Chern roots of F_{n+1-r} are represented by $-y_r, \dots, -y_n$ for each r . When $k = a_1$ we have $\vartheta_r(X; Y_1) = \sum_{i=0}^r q_{r-i}(X) e_i(y_1, \dots, y_{a_1})$ for any $r \geq 0$. Therefore, we obtain

$$\begin{aligned} \pi_n(\vartheta_r(X; Y_1)) &= \sum_{i,j \geq 0} e_{r-i-j}(X) h_j(Y) e_i(-x_1, \dots, -x_{a_1}) \\ &= \sum_{j \geq 0} e_{r-j}(x_1, \dots, x_n, -x_1, \dots, -x_{a_1}) h_j(Y) = c_r(Q_1 - F_n) \end{aligned}$$

and hence $\pi_n(\Theta_\lambda(X; Y_1)) = \Theta_\lambda(Q_1 - F_n)$, for any a_1 -strict partition λ . Moreover, for any partition μ and $r \geq 2$, we have

$$\pi_n(s_\mu(Y_r)) = s_\mu(-x_{a_{r-1}+1}, \dots, -x_{a_r}) = s_\mu(Q_r),$$

while

$$\pi_n(s_\mu(0/Z_r)) = s_{\mu'}(-y_{b_{r-1}+1}, \dots, -y_{b_r}) = s_{\mu'}(\hat{Q}_r) = s_\mu(F_{n+b_{r-1}} - F_{n+b_r}).$$

We deduce that π_n maps formula (24) onto the desired equality. $\square$

5. GIAMBELLI FORMULAS FOR SYMPLECTIC FLAG VARIETIES

5.1. Partial isotropic flag varieties. Equip the vector space $E = \mathbb{C}^{2n}$ with a nondegenerate skew-symmetric bilinear form. Fix a sequence $\mathbf{a} : a_1 < \dots < a_p$ of nonnegative integers with $a_p < n$, and set $a_{p+1} = n$. Let $\mathfrak{X}(\mathbf{a})$ be the variety parametrizing partial flags of subspaces

$$0 = E_0 \subset E_1 \subset \dots \subset E_p \subset E$$

with $\dim E_i = n - a_{p+1-i}$ and E_p isotropic. The same notation will be used to denote the tautological partial flag $E_\bullet$ of vector bundles over $\mathfrak{X}(\mathbf{a})$.

A presentation of the cohomology ring of $\mathfrak{X}(\mathbf{a})$ as a quotient of the symmetric algebra on the characters of a maximal torus in Sp_{2n} is well known [Bo]. We will give here an alternative presentation using the Chern classes of the tautological vector bundles over $\mathfrak{X}(\mathbf{a})$. Let $Q_1 = E/E_p, Q_2 = E_p/E_{p-1}, \dots, Q_{p+1} = E_1$ and set $\sigma_i = c_i(Q_1)$ for $1 \leq i \leq n + a_1$ and $c_j^r = c_j(Q_r)$ for $2 \leq r \leq p + 1$ and $1 \leq j \leq a_r - a_{r-1}$. We then have the following result

Proposition 3. *The cohomology ring $H^*(\mathfrak{X}(\mathbf{a}), \mathbb{Z})$ is presented as a quotient of the polynomial ring $\mathbb{Z}[\sigma_1, \dots, \sigma_{n+a_1}, c_1^2, \dots, c_{a_2-a_1}^2, \dots, c_1^{p+1}, \dots, c_{n-a_p}^{p+1}]$ modulo the relations*

$$\det(\sigma_{1+j-i})_{1 \leq i, j \leq r} = (-1)^r \sum_{i_2 + \dots + i_{p+1} = r} c_{i_2}^2 \dots c_{i_{p+1}}^{p+1}, \quad 1 \leq r \leq n + a_1$$

and

$$(32) \quad \sigma_r^2 + 2 \sum_{i=1}^{n+a_1-r} (-1)^i \sigma_{r+i} \sigma_{r-i} = 0, \quad a_1 + 1 \leq r \leq n.$$

Proof. Let $\mathrm{IG} = \mathrm{IG}(n - a_1, 2n)$ be the Grassmannian parametrizing isotropic subspaces of E of dimension $n - a_1$. According to [BKT1, Thm. 1.2], the cohomology ring of IG is isomorphic to the polynomial ring generated by the Chern classes $\sigma_i = c_i(Q_1)$ and the Chern classes of E_p , modulo the relations

$$\det(\sigma_{1+j-i})_{1 \leq i, j \leq r} = (-1)^r c_r(E_p), \quad 1 \leq r \leq n + a_1$$

coming from the Whitney sum formula $c(E_p)c(Q_1) = 1$, as well as the relations (32). The map $\mathfrak{X}(\mathbf{a}) \rightarrow \mathrm{IG}$ sending $E_\bullet$ to E_p realizes $\mathfrak{X}(\mathbf{a})$ as a fiber bundle over IG with fiber a partial SL_{n-a_1} flag variety. We deduce using e.g. [Gr, Thm. 1] that $H^*(\mathfrak{X}(\mathbf{a}))$ is isomorphic to the polynomial ring $H^*(\mathrm{IG})[c_1^2, \dots, c_{a_2-a_1}^2, \dots, c_1^{p+1}, \dots, c_{n-a_p}^{p+1}]$ modulo the relations

$$\sum_{i_2 + \dots + i_{p+1} = r} c_{i_2}^2 \dots c_{i_{p+1}}^{p+1} = c_r(E_p), \quad 1 \leq r \leq n - a_1.$$

The proposition follows by combining these two facts. $\square$

5.2. Giambelli formulas. Our choice of the special Schubert classes on isotropic Grassmannians agrees with the conventions in [BKT1]. Let $F_\bullet$ be a fixed complete isotropic flag of subspaces in $\mathbb{C}^{2n}$. For each $w \in W^{\mathbf{a}}$, the Schubert variety $\mathfrak{X}_w(F_\bullet)$ in $\mathfrak{X}(\mathbf{a})$ is defined by the same equation (30) as before. Let $E'_j = E/E_{p+1-j}$ for $1 \leq j \leq p$. The Chern classes $c_i(E'_j)$ for all i, j are the Schubert classes on $\mathfrak{X}(\mathbf{a})$ which are pullbacks of special Schubert classes on symplectic Grassmannians. By

definition, they are the *special Schubert classes* on $\mathfrak{X}(\mathfrak{a})$, and they generate the cohomology ring $H^*(\mathfrak{X}(\mathfrak{a}))$ by Proposition 3. Specializing Theorem 3 to the case when the base B is a point gives the next result.

Corollary 4. *For every $w \in W^{\mathfrak{a}}$, we have*

$$\begin{aligned} [\mathfrak{X}_w] &= \sum_{\underline{\lambda}} e_{\underline{\lambda}}^w \Theta_{\lambda^1}(Q_1) s_{\lambda^2}(Q_2) \cdots s_{\lambda^p}(Q_p) \\ &= \sum_{\underline{\lambda}} e_{\underline{\lambda}}^w \Theta_{\lambda^1}(E'_1) s_{\lambda^2}(E'_2 - E'_1) \cdots s_{\lambda^p}(E'_p - E'_{p-1}) \end{aligned}$$

in $H^*(\mathfrak{X}(\mathfrak{a}))$, where the sums are over all sequences of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^p)$ with λ^1 a_1 -strict, and the coefficients $e_{\underline{\lambda}}^w$ are given by (22).

6. SPLITTING ORTHOGONAL SCHUBERT POLYNOMIALS

In this final section we discuss the form of our splitting results for the orthogonal groups; we will work throughout with coefficients in the ring $\mathbb{Z}[\frac{1}{2}]$. For $w \in W_{\infty}$, let $s(w)$ denote the number of i such that $w(i) < 0$. It follows e.g. from [BH, IMN] that the polynomials $\mathfrak{B}_w = 2^{-s(w)} \mathfrak{C}_w$ represent the Schubert classes in the (equivariant) cohomology ring of odd orthogonal flag varieties. Therefore the solutions to the Schubert polynomial splitting and Giambelli problems for types B and C are essentially the same. We will describe the splitting theorems for the even orthogonal groups below; the story is entirely analogous to the symplectic case.

The elements of the Weyl group $\widetilde{W}_n$ for the root system D_n may be represented by signed permutations, as in e.g. [B, KT1]. The group $\widetilde{W}_n$ is an extension of S_n by an element s_0 which acts on the right by

$$(u_1, u_2, \dots, u_n) s_0 = (\overline{u}_2, \overline{u}_1, u_3, \dots, u_n).$$

Let $\widetilde{W}_{\infty} = \cup_n \widetilde{W}_n$ and $\mathbb{N} = \{0, 1, 2, \dots\}$. A *reduced word* of $w \in \widetilde{W}_{\infty}$ is a sequence $a_1 \cdots a_{\ell}$ of elements in $\mathbb{N}$ such that $w = s_{a_1} \cdots s_{a_{\ell}}$ and $\ell = \ell(w)$. We say that w has *descent* at position $r \geq 0$ if $\ell(ws_r) < \ell(w)$, where s_r is the simple reflection indexed by r . If $k \geq 2$, we say that an element $w \in \widetilde{W}_{\infty}$ is *increasing up to k* if it has no descents less than k ; this means that $|w_1| < w_2 < \cdots < w_k$. We agree that all elements of $\widetilde{W}_{\infty}$ are increasing up to 0 and increasing up to 1.

For $k \in \mathbb{N} \setminus \{1\}$, an element $w \in \widetilde{W}_{\infty}$ is *k -Grassmannian* if $\ell(ws_i) = \ell(w) + 1$ for all $i \neq k$. We say that w is *1-Grassmannian* if $\ell(ws_i) = \ell(w) + 1$ for all $i \geq 2$. Following [BKT1, §4], to any k -strict partition λ we associate a number in $\{0, 1, 2\}$ called the *type* of λ , and denoted $\text{type}(\lambda)$. Here ‘type’ is a multi-valued function such that $\text{type}(\lambda) = 0$ if λ has no part equal to k , and $\text{type}(\lambda) \in \{1, 2\}$, otherwise. The geometric significance of the type of λ is explained in [BKT1, §4.5].

Given a k -Grassmannian element $w \in \widetilde{W}_n$, there exist unique strict partitions u, ζ, v of lengths k, r , and $n - k - r$, respectively, so that

$$w = (\widehat{u}_k, \dots, u_1, \overline{\zeta}_1, \dots, \overline{\zeta}_r, v_{n-k-r}, \dots, v_1)$$

where $\widehat{u}_k$ is equal to u_k or $\overline{u}_k$, according to the parity of r . If $\mu_i = u_i + i - k - 1 + \#\{j \mid \zeta_j > u_i\}$, then w corresponds to a typed k -strict partition λ such that the lengths of the first k columns of λ are given by $\mu_1, \dots, \mu_k$. The part of λ in columns $k+1$ and higher is given by $(\zeta_1 - 1, \dots, \zeta_r - 1)$; here it is possible that $\zeta_r = 1$, so that the sequence ends with a zero. Finally, if $\text{type}(\lambda) > 0$, then $\widehat{u}_k$ is unbarred

if and only if $\text{type}(\lambda) = 1$. This defines a bijection between the k -Grassmannian elements of $\widetilde{W}_\infty$ and the k -strict partitions of all three possible types; we let w_λ denote the element of $\widetilde{W}_\infty$ associated to the typed k -strict partition λ .

Following [L1], we will use the nilCoxeter algebra $\widetilde{W}_n$ of $\widetilde{W}_n$ to define type D Stanley symmetric functions. $\widetilde{W}_n$ is the free associative algebra with unity generated by the elements $u_0, u_1, \dots, u_{n-1}$ modulo the relations

$$\begin{aligned} u_i^2 &= 0 & i \geq 0 ; \\ u_0 u_1 &= u_1 u_0 \\ u_0 u_2 u_0 &= u_2 u_0 u_2 \\ u_i u_{i+1} u_i &= u_{i+1} u_i u_{i+1} & i > 0 ; \\ u_i u_j &= u_j u_i & j > i + 1, \text{ and } (i, j) \neq (0, 2). \end{aligned}$$

For any $w \in \widetilde{W}_n$, choose a reduced word $a_1 \cdots a_\ell$ for w and define $u_w = u_{a_1} \cdots u_{a_\ell}$. We denote the coefficient of $u_w \in \widetilde{W}_n$ in the expansion of the element $f \in \widetilde{W}_n$ by $\langle f, w \rangle$. Let t be an indeterminate and, following [L1, 4.4], define

$$D(t) = (1 + tu_{n-1}) \cdots (1 + tu_2)(1 + tu_1)(1 + tu_0)(1 + tu_2) \cdots (1 + tu_{n-1}).$$

According to [L1, Lemma 4.24], we have $D(s)D(t) = D(t)D(s)$ for any commuting variables s, t . If $D(X) = D(x_1)D(x_2) \cdots$, then the functions $E_w(X)$ defined by

$$E_w(X) = \langle D(X), w \rangle$$

are the type D Stanley symmetric functions, in agreement with [BH, §3].

Next, define

$$\mathfrak{D}_w(X; Y, Z) = \left\langle \tilde{A}_{n-1}(z_{n-1}) \cdots \tilde{A}_1(z_1) D(X) A_1(y_1) \cdots A_{n-1}(y_{n-1}), w \right\rangle.$$

The polynomials $\mathfrak{D}_w(X; Y) := \mathfrak{D}_w(X; Y, 0)$ are the type D Billey-Haiman Schubert polynomials, and the $\mathfrak{D}_w(X; Y, Z)$ are their double versions studied in [IMN]. If $w = w_\lambda$ is k -Grassmannian, then $\mathfrak{D}_{w_\lambda}(X; Y)$ is an *eta polynomial* $H_\lambda(X; Y)$; the H_λ are defined using raising operator expansions analogous to (3) in [BKT4]. When $k = 0$, we have that λ is a strict partition and $H_\lambda(X; Y) = E_{w_\lambda}(X) = P_\lambda(X)$ is a Schur P -function.

Definition 4. Given $w \in \widetilde{W}_\infty$, the *type D mixed Stanley function* $I_w(X; Y)$ is defined by the equation

$$I_w(X; Y) = \langle D(X)A(Y), w \rangle = \sum_{uv=w} E_u(X)G_v(Y)$$

summed over all reduced factorizations $uv = w$ with $v \in S_\infty$.

One checks that if w is increasing up to k , then $I_w^{(k)} = \mathfrak{D}_w^{(k)}(X; Y)$; in particular, if $w = w_\lambda$ is k -Grassmannian, then $I_{w_\lambda}^{(k)} = H_\lambda$. Furthermore, it follows from [B, Thms. 4, 5] that the I_w satisfy the type D transition equations

$$I_w^{(k)} = \sum_{\substack{1 \leq i < r \\ \ell(w t_{rs} t_{ir}) = \ell(w)}} I_{w t_{rs} t_{ir}}^{(k)} + \sum_{\substack{i \neq r \\ \ell(w t_{rs} \bar{t}_{ir}) = \ell(w)}} I_{w t_{rs} \bar{t}_{ir}}^{(k)}$$

where r is the last positive descent of w and s is maximal such that $w_s < w_r$.

For any $w \in \widetilde{W}_\infty$ which is increasing up to k , we construct the k -transition tree $\tilde{T}^k(w)$ with nodes given by elements of $\widetilde{W}_\infty$ and root w as in §2.2. Let r be the last descent of w . If $w = 1$, or $k \neq 1$ and $r = k$, or $k = 1$ and $r \in \{0, 1\}$, then set

$\tilde{T}^k(w) = \{w\}$. Otherwise, let $s = \max(i > r \mid w_i < w_r)$ and $\tilde{\Phi}(w) = \tilde{\Phi}_1(w) \cup \tilde{\Phi}_2(w)$, where

$$\begin{aligned}\tilde{\Phi}_1(w) &= \{wt_{rs}t_{ir} \mid 1 \leq i < r \text{ and } \ell(wt_{rs}t_{ir}) = \ell(w)\}, \\ \tilde{\Phi}_2(w) &= \{wt_{rs}\bar{t}_{ir} \mid i \neq r \text{ and } \ell(wt_{rs}\bar{t}_{ir}) = \ell(w)\}.\end{aligned}$$

To define $\tilde{T}^k(w)$, we join w by an edge to each $v \in \tilde{\Phi}(w)$, and attach to each $v \in \tilde{\Phi}(w)$ its tree $\tilde{T}^k(v)$.

The assertions of Lemma 1 remain true for $\tilde{T}^k(w)$, with similar proof. When $k = 0$, this is contained in [B, Thm. 4]. For the case when $r > k > 1$, $\tilde{\Phi}_1(w) = \emptyset$, and $w_s > 0$, one observes that $wt_{rs}\bar{t}_{1r} \in \tilde{\Phi}_2(w)$. We deduce that for any $w \in \widetilde{W}_\infty$ which is increasing up to k ,

$$(33) \quad \mathfrak{D}_w^{(k)} = I_w^{(k)} = \sum_{\lambda: |\lambda| = \ell(w)} d_\lambda^w H_\lambda$$

where the sum is over typed k -strict partitions λ and d_λ^w denotes the number of leaves of $\tilde{T}^k(w)$ of shape λ . Moreover, for any such w , we have

$$(34) \quad \mathfrak{D}_w(X; Y) = \sum_{u(1_k \times v) = w} I_u^{(k)}(X; Y) \mathfrak{S}_v(Y_{>k})$$

where the sum is over all reduced factorizations $u(1_k \times v) = w$ with $v \in S_\infty$.

Using equations (33) and (34), we obtain splitting theorems for the single and double type D Schubert polynomials $\mathfrak{D}_w$, as in §3. Fix two sequences $\mathbf{a} : a_1 < \dots < a_p$ and $\mathbf{b} : 0 = b_1 < \dots < b_q$ of nonnegative integers and set $Y_i = \{y_{a_{i-1}+1}, \dots, y_{a_i}\}$ and $Z_j = \{z_{b_{j-1}+1}, \dots, z_{b_j}\}$ for each i, j .

Theorem 4. *Suppose that $w \in \widetilde{W}_\infty$ is compatible with the sequence $\mathbf{a}$. Then we have*

$$\mathfrak{D}_w(X; Y) = \sum_{u_1 \cdots u_p = w} I_{u_1}(X; Y_1) G_{u_2}(Y_2) \cdots G_{u_p}(Y_p)$$

summed over all reduced factorizations $u_1 \cdots u_p = w$ compatible with $\mathbf{a}$ such that $u_2, \dots, u_p \in S_\infty$.

Given any sequence of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^{p+q-1})$ with λ^q a_1 -strict and typed, we define

$$g_{\underline{\lambda}}^w = \sum_{u_1 \cdots u_{p+q-1} = w} c_{\lambda^1}^{u_1} \cdots c_{\lambda^{q-1}}^{u_{q-1}} d_{\lambda^q}^{u_q} c_{\lambda^{q+1}}^{u_{q+1}} \cdots c_{\lambda^{p+q-1}}^{u_{p+q-1}},$$

where the sum is over reduced factorizations $u_1 \cdots u_{p+q-1} = w$ compatible with $\mathbf{a}$, $\mathbf{b}$ such that $u_i \in S_\infty$ for all $i \neq q$, and the integers $c_{\lambda^i}^{u_i}$ and $d_{\lambda^q}^{u_q}$ are as in (4) and (33), respectively.

Corollary 5. *Suppose that w and w^{-1} are compatible with the sequences $\mathbf{a}$ and $\mathbf{b}$, respectively. Then $\mathfrak{D}_w(X; Y, Z)$ is equal to*

$$\sum_{u_1 \cdots u_{p+q-1} = w} G_{u_1}(0/Z_q) \cdots G_{u_{q-1}}(0/Z_2) I_{u_q}(X; Y_1) G_{u_{q+1}}(Y_2) \cdots G_{u_{p+q-1}}(Y_p)$$

TABLE 1. Schubert polynomials with w increasing up to 1

w	word	$\mathfrak{C}_w(X; Y)$	w	word	$\mathfrak{D}_w(X; Y)$
123		1	123		1
213	1	Θ_1	213	1	H_1
132	2	$\Theta_1 + y_2$	$\overline{213}$	0	H'_1
$\overline{213}$	01	Θ_2	132	2	$H_1 + H'_1 + y_2$
312	21	$\Theta_{(1,1)}$	$\overline{123}$	01	H_2
231	12	$\Theta_2 + \Theta_1 y_2$	312	21	$H_{(1,1)}$
$\overline{123}$	101	Θ_3	$\overline{312}$	20	$H'_{(1,1)}$
$\overline{312}$	021	$\Theta_{(2,1)}$	231	12	$H_2 + H_1 y_2$
321	212	$\Theta_{(2,1)} + \Theta_{(1,1)} y_2$	$\overline{231}$	02	$H_2 + H'_1 y_2$
$\overline{231}$	012	$\Theta_3 + \Theta_2 y_2$	$\overline{132}$	201	H_3
$\overline{132}$	2101	Θ_4	$\overline{321}$	021	$H_{(2,1)}$
$\overline{321}$	1021	$\Theta_{(3,1)}$	$\overline{321}$	120	$H'_{(2,1)}$
$\overline{132}$	1012	$\Theta_4 + \Theta_3 y_2$	$\overline{132}$	012	$H_3 + H_2 y_2$
$\overline{321}$	0212	$\Theta_4 + \Theta_{(3,1)} + \Theta_{(2,1)} y_2$	321	212	$H_3 + H_{(2,1)} + H_{(1,1)} y_2$
$\overline{231}$	21021	$\Theta_{(4,1)}$	$\overline{321}$	202	$H_3 + H'_{(2,1)} + H'_{(1,1)} y_2$
$\overline{321}$	01021	$\Theta_{(3,2)}$	$\overline{231}$	2021	$H_{(3,1)}$
$\overline{123}$	21012	$\Theta_5 + \Theta_4 y_2$	$\overline{231}$	2120	$H'_{(3,1)}$
$\overline{312}$	10212	$\Theta_{(4,1)} + \Theta_{(3,1)} y_2$	$\overline{123}$	2012	$H_4 + H_3 y_2$
$\overline{231}$	021021	$\Theta_{(4,2)}$	$\overline{312}$	0212	$H_{(3,1)} + H_{(2,1)} y_2$
213	210212	$\Theta_{(5,1)} + \Theta_{(4,1)} y_2$	$\overline{312}$	1202	$H'_{(3,1)} + H'_{(2,1)} y_2$
$\overline{312}$	010212	$\Theta_{(4,2)} + \Theta_{(3,2)} y_2$	$\overline{132}$	12021	$H_{(3,2)}$
$\overline{132}$	1021021	$\Theta_{(4,3)}$	$\overline{213}$	20212	$H_{(4,1)} + H_{(3,1)} y_2$
$\overline{213}$	0210212	$\Theta_{(5,2)} + \Theta_{(4,2)} y_2$	$\overline{213}$	21202	$H'_{(4,1)} + H'_{(3,1)} y_2$
$\overline{123}$	10210212	$\Theta_{(5,3)} + \Theta_{(4,3)} y_2$	$\overline{123}$	120212	$H_{(4,2)} + H_{(3,2)} y_2$

summed over all reduced factorizations $u_1 \cdots u_{p+q-1} = w$ compatible with $\mathbf{a}, \mathbf{b}$ such that $u_i \in S_\infty$ for all $i \neq q$. Furthermore, we have

$$(35) \quad \mathfrak{D}_w(X; Y, Z) = \sum_{\underline{\lambda}} g_{\underline{\lambda}}^w s_{\lambda^1}(0/Z_q) \cdots H_{\lambda^q}(X; Y_1) \cdots s_{\lambda^{p+q-1}}(Y_p)$$

summed over all sequences of partitions $\underline{\lambda} = (\lambda^1, \dots, \lambda^{p+q-1})$ with λ^q a_1 -strict and typed.

In the same manner as for the symplectic groups, the above splitting results imply Giambelli and degeneracy locus formulas for the orthogonal groups. We use the Giambelli formula for even orthogonal Grassmannians from [BKT4] in (35).

Table 1 lists the Billey-Haiman Schubert polynomials for the root systems of type C_3 and D_3 indexed by the elements w in the respective Weyl groups which are increasing up to 1. In each case, the polynomial is written as a positive sum of $k = 1$ theta and eta polynomials in the variables (X, y_1) times $s_j(y_2)$ for $j \in \{0, 1\}$. The primed eta polynomials H'_λ are indexed by 1-strict partitions λ of type 2.

REFERENCES

- [BGG] I. N. Bernstein, I. M. Gelfand and S. I. Gelfand : *Schubert cells and cohomology of the spaces G/P* , Russian Math. Surveys **28** (1973), 1–26.
- [Be] A. Bertram : *Quantum Schubert calculus*, Adv. Math. **128** (1997), 289–305.
- [B] S. Billey : *Transition equations for isotropic flag manifolds*, Discrete Math. **193** (1998), 69–84.

- [BJS] S. Billey, W. Jockusch and R. P. Stanley : *Some combinatorial properties of Schubert polynomials*, J. Algebraic Combin. **2** (1993), 345–374.
- [BH] S. Billey and M. Haiman : *Schubert polynomials for the classical groups*, J. Amer. Math. Soc. **8** (1995), 443–482.
- [BL] S. Billey and T. K. Lam : *Vexillary elements in the hyperoctahedral group*, J. Algebraic Combin. **8** (1998), 139–152.
- [Bo] A. Borel : *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math. **57** (1953), 115–207.
- [BKT1] A. S. Buch, A. Kresch and H. Tamvakis : *Quantum Pieri rules for isotropic Grassmannians*, Invent. Math., to appear.
- [BKT2] A. S. Buch, A. Kresch and H. Tamvakis : *A Giambelli formula for isotropic Grassmannians*, Preprint (2008), available at arXiv:0811.2781.
- [BKT3] A. S. Buch, A. Kresch and H. Tamvakis : *Quantum Giambelli formulas for isotropic Grassmannians*, Preprint (2008), available at arXiv:0812.0970.
- [BKT4] A. S. Buch, A. Kresch and H. Tamvakis : *A Giambelli formula for even orthogonal Grassmannians*, in preparation.
- [BKTY] A. S. Buch, A. Kresch, H. Tamvakis, and A. Yong : *Schubert polynomials and quiver formulas*, Duke Math. J. **122** (2004), 125–143.
- [BF] A. S. Buch and W. Fulton : *Chern class formulas for quiver varieties*, Invent. Math. **135** (1999), 665–687.
- [CF] I. Ciocan-Fontanine : *On quantum cohomology rings of partial flag varieties*, Duke Math. J. **98** (1999), 485–524.
- [D1] M. Demazure : *Invariants symétriques des groupes de Weyl et torsion*, Invent. Math. **21** (1973), 287–301.
- [D2] M. Demazure : *Désingularisation des variétés de Schubert généralisées*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **7** (1974), 53–88.
- [EG] M. Edelman and C. Greene : *Balanced tableaux*, Adv. Math. **63** (1987), 42–99.
- [FGP] S. Fomin, S. Gelfand and A. Postnikov : *Quantum Schubert polynomials*, J. Amer. Math. Soc. **10** (1997), 565–596.
- [FG] S. Fomin and C. Greene : *Noncommutative Schur functions and their applications*, Discrete Math. **193** (1998), 179–200.
- [FK1] S. Fomin and A. N. Kirillov : *The Yang-Baxter equation, symmetric functions, and Schubert polynomials*, Discrete Math. **153** (1996), 123–143.
- [FK2] S. Fomin and A. N. Kirillov : *Combinatorial B_n -analogs of Schubert polynomials*, Trans. Amer. Math. Soc. **348** (1996), 3591–3620.
- [FS] S. Fomin and R. P. Stanley : *Schubert polynomials and the nil-Coxeter algebra*, Adv. in Math. **103** (1994), 196–207.
- [Fu1] W. Fulton : *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), 381–420.
- [Fu2] W. Fulton : *Schubert varieties in flag bundles for the classical groups*, Proceedings of the Hirzebruch 65 Conference on Algebraic Geometry (Ramat Gan, 1993), 241–262, Israel Math. Conf. Proc. **9**, Ramat Gan, 1996.
- [Fu3] W. Fulton : *Determinantal formulas for orthogonal and symplectic degeneracy loci*, J. Differential Geom. **43** (1996), 276–290.
- [Fu4] W. Fulton : *Universal Schubert polynomials*, Duke Math. J. **96** (1999), 575–594.
- [FP] W. Fulton and P. Pragacz : *Schubert varieties and degeneracy loci*, Lecture Notes in Math. **1689**, Springer-Verlag, Berlin, 1998.
- [G] G. Z. Giambelli : *Risoluzione del problema degli spazi secanti*, Mem. R. Accad. Sci. Torino (2) **52** (1902), 171–211.
- [Gra] W. Graham : *The class of the diagonal in flag bundles*, J. Differential Geom. **45** (1997), 471–487.
- [Gr] A. Grothendieck : *Quelques propriétés fondamentales en théorie des intersections*, Séminaire C. Chevalley **3** (1958), exp. no. 4, 1–36.
- [H] M. D. Haiman : *Dual equivalence with applications, including a conjecture of Proctor*, Discrete Math. **99** (1992), 79–113.
- [IMN] T. Ikeda, L. Mihalcea, and H. Naruse : *Double Schubert polynomials for the classical groups*, Preprint (2008), arXiv:0810.1348.

- [KMS] A. Knutson, E. Miller, and M. Shimozono : *Four positive formulae for type A quiver polynomials*, Invent. Math. **166** (2006), 229–325.
- [KL] G. Kempf and D. Laksov : *The determinantal formula of Schubert calculus*, Acta Math. **132** (1974), 153–162.
- [Kr] W. Krawkiewicz : *Reduced decompositions in hyperoctahedral groups*, C. R. Acad. Sci. Paris Sér. I Math. **309** (1989), 903–907.
- [KT1] A. Kresch and H. Tamvakis : *Double Schubert polynomials and degeneracy loci for the classical groups*, Ann. Inst. Fourier **52** (2002), 1681–1727.
- [KT2] A. Kresch and H. Tamvakis : *Quantum cohomology of the Lagrangian Grassmannian*, J. Algebraic Geom. **12** (2003), 777–810.
- [KT3] A. Kresch and H. Tamvakis : *Quantum cohomology of orthogonal Grassmannians*, Compos. Math. **140** (2004), 482–500.
- [L1] T. K. Lam : *B and D analogues of stable Schubert polynomials and related insertion algorithms*, Ph.D. thesis, M.I.T., 1994.
- [L2] T. K. Lam : *B_n Stanley symmetric functions*, Discrete Math. **157** (1996), 241–270.
- [LP1] A. Lascoux and P. Pragacz : *Operator calculus for $\tilde{Q}$ -polynomials and Schubert polynomials*, Adv. Math. **140** (1998), 1–43.
- [LP2] A. Lascoux and P. Pragacz : *Orthogonal divided differences and Schubert polynomials, $\tilde{P}$ -functions, and vertex operators*, Michigan Math. J. **48** (2000), 417–441.
- [LS1] A. Lascoux and M.-P. Schützenberger : *Polynômes de Schubert*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982), 447–450.
- [LS2] A. Lascoux and M.-P. Schützenberger : *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), 629–633.
- [LS3] A. Lascoux and M.-P. Schützenberger : *Schubert polynomials and the Littlewood-Richardson rule*, Lett. Math. Phys. **10** (1985), 111–124.
- [M1] I. G. Macdonald : *Notes on Schubert polynomials*, Publ. LACIM **6**, Univ. de Québec à Montréal, Montréal, 1991.
- [M2] I. Macdonald : *Symmetric Functions and Hall Polynomials*, Second edition, Clarendon Press, Oxford, 1995.
- [Ma] L. Manivel : *Fonctions symétriques, polynômes de Schubert et lieux de dégénérescence*, Cours Spécialisés, 3, Société Mathématique de France, Paris, 1998.
- [Mo] D. Monk : *The geometry of flag manifolds*, Proc. London Math. Soc. **3** (1959), 253–286.
- [P] P. Pragacz : *Algebro-geometric applications of Schur S- and Q-polynomials*, Séminaire d’Algèbre Dubreil-Malliavin 1989-1990, Lecture Notes in Math. **1478** (1991), 130–191, Springer-Verlag, Berlin, 1991.
- [PR] P. Pragacz and J. Ratajski : *Formulas for Lagrangian and orthogonal degeneracy loci; $\tilde{Q}$ -polynomial approach*, Compositio Math. **107** (1997), 11–87.
- [Sa] B. Sagan : *Shifted tableaux, Schur Q-functions, and a conjecture of R. Stanley*, J. Combin. Theory Ser. A **45** (1987), 62–103.
- [Sch] I. Schur : *Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene lineare Substitutionen*, J. reine angew. Math. **139** (1911), 155–250.
- [Sh] M. Shimozono : *Multiplying Schur Q-functions*, J. Combin. Theory Ser. A **87** (1999), 198–232.
- [Sta] R. P. Stanley : *On the number of reduced decompositions of elements of Coxeter groups*, European J. Combin. **5** (1984), 359–372.
- [St1] J. R. Stembridge : *Shifted tableaux and the projective representations of symmetric groups*, Adv. Math. **74** (1989), 87–134.
- [St2] J. R. Stembridge : *Some combinatorial aspects of reduced words in finite Coxeter groups*, Trans. Amer. Math. Soc. **349** (1997), 1285–1332.
- [T1] H. Tamvakis : *Schubert polynomials and Arakelov theory of symplectic flag varieties*, Preprint (2008), arXiv:0808.1329.
- [T2] H. Tamvakis : *Giambelli, Pieri, and tableau formulas via raising operators*, Preprint (2008), arXiv:0812.0639.
- [T3] H. Tamvakis : *Schubert polynomials and Arakelov theory of orthogonal flag varieties*, Preprint (2009), arXiv:0907.3308.
- [Tu] L. W. Tu : *Degeneracy loci*, Proc. conf. algebraic geom. (Berlin, 1985), 296–305, Teubner-Texte Math. **92**, Teubner, Leipzig, 1986.

- [W] D. R. Worley : *A theory of shifted Young tableaux*, Ph.D. thesis, MIT, 1984.
- [Yo] A. Yong : *On combinatorics of quiver component formulas*, J. Algebraic Combin. **21** (2005), 351–371.
- [Y] A. Young : *On quantitative substitutional analysis VI*, Proc. Lond. Math. Soc. (2) **34** (1932), 196–230.

UNIVERSITY OF MARYLAND, DEPARTMENT OF MATHEMATICS, 1301 MATHEMATICS BUILDING,
COLLEGE PARK, MD 20742, USA

E-mail address: `harryt@math.umd.edu`